

Comments on the Paper by D. G. Rea and B. T. O'Leary, 'On the Composition of the Venus Clouds'

JAMES E. HANSEN AND HOWARD CHEYNEY

*Goddard Space Flight Center, Institute for Space Studies, NASA
New York, New York 10025*

In a recent paper *Rea and O'Leary* [1968] examined the cases for and against ice clouds on Venus and concluded that 'there is no direct evidence for H_2O ice crystals, and furthermore, if they do exist on Venus and dominate the near-infrared scattering process, they must be submicron in diameter.' We believe that at least the latter of these two conclusions is not correct, because clouds composed of ice particles a few microns in diameter do not generally cause absorption features at 1.5 and 2.0 μ as strong as assumed by Rea and O'Leary. Their conclusion was based in part on laboratory reflectivity spectra obtained by *Zander* [1966], but we believe that *Zander's* reflectivity spectra are representative of particles larger by a factor of about 5 than suggested in *Zander's* publication.

Rea and O'Leary examined spectra of Venus obtained by *Kuiper* [1962] with an earth-based telescope for wavelengths $1.4 \mu \lesssim \lambda \lesssim 2.3 \mu$ and balloon observations of *Bottima et al.* [1964, 1965] for $1.8 \mu \lesssim \lambda \lesssim 3.4 \mu$. They showed that the absorption near 1.5 and 2.0 μ is due largely to CO_2 ; hence, except for the reflectivity minimum near 3 μ , there is no significant evidence for the broad-band absorption features that would be expected for ice in the near-infrared. Rea and O'Leary illustrated spectral reflectivities of ice clouds having particle diameters $\approx 10\text{--}100 \mu$, and these showed very strong absorption features at 1.5 and 2.0 μ . Although the strength of these features would diminish for smaller particles, it was shown that the features were nearly as strong in laboratory measurements by *Zander* [1966] in which the particles were reported as having diameters primarily in the range $1 \mu \leq d \leq 3 \mu$. From this they argued that either ice particles are not the principal scatterers of the infrared radiation on Venus or, if they are, the particle

diameters must be less than 1 μ and probably less than 0.1 μ . Since the planet-wide existence of submicron ice particles is unlikely, they concluded that the major scatterers of radiation on Venus are almost certainly not composed of H_2O ice.

However, solutions of the radiative transfer problem for diffuse reflection from ice clouds have been made [*Hansen and Cheyney*, 1968] for a size distribution of spherical particles and compared with *Zander's* observations. The optical constants for ice were taken from the tables compiled by *Irvine and Pollack* [1968]; from these the absorption and scattering cross sections and the phase function were calculated for each particle, weighted for the mixture, and employed in the transfer problem to obtain the complete angular distribution of reflected light. When the size distribution specified in *Zander's* article (diameters of 1–3 μ) was employed, the absorption features at 1.5 and 2.0 μ were found to be much too weak to be compatible with the laboratory observations; however, it was learned from *Zander* (private communication, 1968) that a significant number of the 1–3 μ particles were clustered into tightly packed (void space $< 20\%$) aggregates as large as 15 μ . Since the components of an aggregate do not scatter incoherently, it is a better approximation to treat each aggregate as a single large particle, and a relatively small number of these may completely change the scattering properties of a unit volume. According to *Zander* (private communication, 1968) approximately 20% of the particles were aggregates (with an aggregate counted as being one particle) and the distribution of the aggregates peaked at $d \approx 10 \mu$. When the complete distribution including aggregates was used in the radiative transfer problem, a close fit was obtained with *Zander's* observed curve.

From the apparent discrepancy between Sagan and Pollack's [1967] calculations and Zander's laboratory spectra, Rea and O'Leary suggested, along with other possibilities, that Sagan and Pollack's approximate radiative transfer solution may be in serious error, but our results for a narrow size distribution are compatible with the single particle calculations of Sagan and Pollack. Certainly their two-stream approximation cannot cause an order of magnitude error in the particle sizing.

The primary point that we wish to make is that with micron-sized ice particles as the major scatterers the reflectivity minima near 1.5 and 2.0 μ are not generally as large as assumed by Rea and O'Leary, and hence Kuiper's ground-based observations may be compatible with clouds of such particles. For example, with a particle size distribution having diameters $1 \mu \leq d \leq 3 \mu$, the maximum of the distribution being at 1.7 μ , we obtain the reflectivities in Table 1, where the reflectivity is the ratio of the calculated intensity to the intensity of a Lambert surface having albedo unity, and τ is the optical thickness of the ice clouds at $\lambda = 0.95 \mu$. The computations were made for the angles of incidence and emergence in Zander's experiment and hence the results are not identical with the integrated light from Venus at a given phase angle, but Rea and O'Leary's conclusions were also based

on Zander's experiment. The table illustrates that with a finite cloud of micron-sized ice particles the absorption features at 1.5 and 2.0 μ are weak, and hence the spectroscopic observations must be accurate in order to establish or rule out the existence of ice clouds.

We do not maintain that this represents positive evidence for ice clouds on Venus; we do believe, however, that the arguments against having clouds of micron-sized ice crystals are not yet conclusive.

Acknowledgments. We thank Dr. R. Jastrow, director of the Institute for Space Studies, for his hospitality and provision of excellent facilities; Drs. J. Pollack, W. Plummer, D. Rea, and P. Thaddeus for helpful discussions; and Dr. R. Zander for valuable correspondence.

During the course of this research J. E. H. held a National Research Council Postdoctoral Research Associateship supported by the National Aeronautics and Space Administration and H. C. was supported by grant NsG 399/33-016-013 through the Department of Meteorology and Oceanography, New York University.

REFERENCES

- Bottema, M., W. Plummer, J. Strong, and R. Zander, Composition of the clouds of Venus, *Astrophys. J.*, **140**, 1640, 1964.
- Bottema, M., W. Plummer, J. Strong, and R. Zander, The composition of the Venus clouds and implications for model atmospheres, *J. Geophys. Res.*, **70**, 4401, 1965.
- Hansen, J., and H. Cheyney, Theoretical spectral scattering of ice clouds in the near infrared, to be published, 1968.
- Irvine, W. M., and J. B. Pollack, Infrared optical properties of water and ice spheres, *Icarus*, **8**, 324, 1968.
- Kuiper, G. P., Photometry of the infrared spectrum of Venus, 1-2.5 microns, *Commun. Lunar Planetary Lab.*, **1**, 83, 1962.
- Rea, D. G., and B. T. O'Leary, On the composition of the Venus clouds, *J. Geophys. Res.*, **73**, 665, 1968.
- Sagan, C., and J. B. Pollack, Anisotropic non-conservative scattering and the clouds of Venus, *J. Geophys. Res.*, **72**, 469, 1967.
- Zander, R., Spectral scattering properties of ice clouds and hoarfrost, *J. Geophys. Res.*, **71**, 375, 1966.

TABLE 1. Reflectivity for Particle Size Distribution between 1 and 3 μ

λ, μ	Reflectivity for τ Values of		
	8	16	32
1.52	0.25	0.45	0.60
1.7	0.24	0.46	0.67
1.85	0.23	0.45	0.68
2.0	0.18	0.35	0.47
2.15	0.18	0.37	0.55
3.0	0.003	0.003	0.003
3.2	0.02	0.02	0.02
3.4	0.13	0.14	0.14
Spherical albedo at $\lambda = 1.85 \mu$	40%	56%	70%

(Received February 19, 1968;
revised May 3, 1968.)